

~~CPA - 1951~~

2780 - 6334

Photoacclimation of phytoplankton in different biogeochemical provinces of the Southern Ocean and its significance for estimating primary production

Die Photoakklimatisierung von Phytoplankton in verschiedenen biogeochemischen Provinzen des Antarktischen Ozeans und ihre Bedeutung für die Abschätzung der Primärproduktion

Astrid Bracher

Ber. Polarforsch. 341 (1999)

ISSN 0176 - 5027

TABLE OF CONTENTS

ABBREVIATIONS	III
SUMMARY	V
ZUSAMMENFASSUNG	VIII
1. GENERAL INTRODUCTION	1
1.1 Marine primary production	1
1.2 Southern Ocean	2
1.2.1 Primary production and the Antarctic Paradox	5
1.2.2 Ozone hole and ultraviolet radiation	9
1.3 Methodological considerations	12
1.4 Thesis outline	13
2. THEORETICAL ASPECTS OF MARINE PRIMARY PRODUCTION	16
2.1 Underwater radiation	16
2.2 Photosynthetic pigments	19
2.3 <i>In vivo</i> light absorption	20
2.4 Photosynthetic responses: The P versus E curve	21
2.5 Photoacclimation and photoinhibition	24
3. SUMMARY OF RESULTS	28
4. GENERAL DISCUSSION	32
4.1 Introduction	32
4.2 Biogeochemical provinces within the Southern Ocean	32
4.2.1 Effect of the upper mixed layer on the formation of biogeochemical provinces	34
4.2.2 Effects by iron and silicate availability	37
4.2.3 Phytoplankton communities	38
4.2.4 Contribution of different regions on total biomass of the Southern Ocean	40
4.3 Biooptical characteristics of phytoplankton	41
4.3.1 Total light absorption by phytoplankton	42
4.3.2. Specific absorption by phytoplankton	43
4.3.3 Quantum yield	48
4.3.4 Sensitivity to enhanced UVR	49
4.4 Methodological aspects	54

4.4.1. Spectral resolution of the underwater light field and absorbance characteristics	54
4.4.2. Areal daily primary production rates	56
4.5 Application to remote sensing	60
4.5.1. Obtaining pigment concentration	60
4.5.2. Obtaining primary production rates	65
4.6 Concluding remarks and future perspectives	66
5. REFERENCES	71
DANKSAGUNG	89