

# **The significance of clay minerals in active fault zones**

Habilitationsschrift

zur Erlangung des akademischen Grades doctor rerum naturalium habitatus

(Dr. rer. nat. habil.)

in der Wissenschaftsdisziplin GEOLOGIE

eingereicht am 25.05.2018 an der

Mathematisch-Naturwissenschaftlichen Fakultät der

Universität Potsdam

von

Anja Maria Schleicher

geboren am 09.09.1971 in Ludwigshafen am Rhein

---

# Content

<b>1. Introduction</b>	1
<b>2. State of the art</b>	3
2.1. Fault zone weakening	3
2.2. Clay mineral formation in fault zones	4
2.3. Drilling into fault zones	5
2.4. Study areas and projects	6
2.4.1. The San Andreas Fault and the SAFOD project	6
2.4.2. The Nankai Trough and the NanTroSEIZE project	8
2.4.3. The Japan Trench and the JFAST project	9
2.4.4. The Alpine Fault and the DFDP project	10
2.5. Major identification methods of clay minerals	11
2.5.1. X-ray diffractometry (XRD)	12
2.5.2. X-ray texture goniometry (XTG)	13
2.5.3. Electron microscopy (SEM/TEM)	13
<b>3. Manuscripts published between 2006-2015</b>	15
3.1. Clay minerals in the San Andreas Fault	17
Origin and significance of clay-coated fractures in mudrock fragments of the SAFOD borehole (Parkfield, California)	19
Electron microscopy of clay minerals in mudrocks from the San Andreas Fault Observatory at Depth (SAFOD)	25
Constraints on mineralization, fluid-rock interaction, and mass transfer during faulting at 2-3 km depth from the SAFOD drill hole	29
On the origin of mixed-layered clay minerals from the San Andreas Fault at 2.5-3 km vertical depth (SAFOD drillhole at Parkfield, California)	41
Nanocoatings of clay and creep of the San Andreas Fault at Parkfield, California	57
Chlorite-smectite clay minerals and fault behavior: new evidence from the San Andreas Fault Observatory at Depth (SAFOD) core	61
3.2. Clay minerals in the Nankai Trough and the Japan Trench subduction zones	73
Constraining clay hydration state and its role in active fault systems	75
Response of natural smectite to seismogenic heating and potential implications for the 2011 Tohoku earthquake in the Japan Trench	89
Structural evolution of an inner accretionary wedge and forearc basin initiation, Nankai margin, Japan	93
3.3. Clay minerals in the Alpine Fault, New Zealand	103
Clay mineral formation and fabric development in the DFDP-1B borehole, central Alpine Fault, New Zealand	105
<b>4. Synthesis and significance of the work</b>	115
4.1. Formation of smectitic nano-coatings	115
4.2. Stability of chlorite-smectite clay minerals at depth	117
4.3. Reaction of clay minerals to quick and protracted heat change	118
4.4. Fabric formation in clay gouge	120
4.5. Concluding remarks	121
<b>5. Theses</b>	123
<b>6. Outlook</b>	125

---

<b>Appendix A: Methods and experimental analytics</b>	<b>129</b>
A.1 X-ray diffractometry	129
A.2 Attachment humidity chamber	130
A.3 Attachment high temperature chamber	130
A.4 X-ray texture goniometry	131
A.5 Scanning electron microscopy	133
A.6 Transmission electron microscopy	134
<b>Appendix B: Publications and conference contributions</b>	<b>137</b>
B.1 List of publications	137
B.2 List of conference talks	141
B.3 List of conference posters	142
B.4 Invited talks	145
<b>Appendix C: Teaching and CV</b>	<b>147</b>
<b>Appendix D: Literature</b>	<b>151</b>